



**Concept-Development Practice Page 6-1**


**Friction**




1. A crate filled with delicious junk food rests on a horizontal floor. Only gravity and the support force of the floor act on it, as shown by the vectors for weight **W** and normal force **n**.
  - a. The net force on the crate is (zero) (greater than zero).
  - b. Evidence for this is \_\_\_\_\_.



2. A slight pull **P** is exerted on the crate, not enough to move it.
  - a. The force of friction **f** acting on the crate is (less than) (equal to) (greater than) **P**.
  - b. The net force on the crate is (zero) (greater than zero).



3. Pull **P** is increased until the crate begins to move. It is pulled so that it moves with constant velocity across the floor.
  - a. Friction **f** is (less than) (equal to) (greater than) **P**.
  - b. Constant velocity means acceleration is (zero) (greater than zero).
  - c. The net force on the crate is (less than) (equal to) (greater than) zero.



4. Pull **P** is further increased and is now greater than friction **f**.
  - a. The net force on the crate is (less than) (equal to) (greater than) zero.
  - b. The net force acts toward the right, so acceleration acts toward the (left) (right).

5. If the pulling force **P** is 150 N and the crate doesn't move, what is the magnitude of **f**? \_\_\_\_\_
6. If the pulling force **P** is 200 N and the crate doesn't move, what is the magnitude of **f**? \_\_\_\_\_
7. If the force of sliding friction is 250 N, what force is necessary to keep the crate sliding at constant velocity? \_\_\_\_\_
8. If the mass of the crate is 50 kg and sliding friction is 250 N, what is the acceleration of the crate when the pulling force is 250 N? \_\_\_\_\_ 300 N? \_\_\_\_\_ 500 N? \_\_\_\_\_

**CONCEPTUAL PHYSICS**

## Falling and Air Resistance


Bronco skydives and parachutes from a stationary helicopter. Various stages of fall are shown in positions (a) through (f). Using Newton's second law,

$$a = \frac{F_{NET}}{m} = \frac{W - R}{m}$$

find Bronco's acceleration at each position (answer in the blanks to the right). You need to know that Bronco's mass  $m$  is 100 kg so his weight is a constant 1000 N. Air resistance  $R$  varies with speed and cross-sectional area as shown.

Circle the correct answers.

- When Bronco's speed is least, his acceleration is  
(least) (most).
- In which position(s) does Bronco experience a downward acceleration?  
(a) (b) (c) (d) (e) (f)
- In which position(s) does Bronco experience an upward acceleration?  
(a) (b) (c) (d) (e) (f)
- When Bronco experiences an upward acceleration, his velocity is  
(still downward) (upward also).
- In which position(s) is Bronco's velocity constant?  
(a) (b) (c) (d) (e) (f)
- In which position(s) does Bronco experience terminal velocity?  
(a) (b) (c) (d) (e) (f)
- In which position(s) is terminal velocity greatest?  
(a) (b) (c) (d) (e) (f)
- If Bronco were heavier, his terminal velocity would be  
(greater) (less) (the same).



## CONCEPTUAL PHYSICS